

Exhibit D



National
Library
of Medicine

My NC
[Sign In] [Regis]

Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book

Search PubMed for

Limits Preview/Index History Clipboard Details

About Entrez

Text Version

Entrez PubMed
Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services
Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
My NCBI (Cubby)

Related Resources
Order Documents
NLM Catalog
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

1: Rev Neurol. 1996 Dec;24(136):1507-12.

Related Articles, Li

[Investigation of corticospinal dysfunction in multiple sclerosis and motoneuron diseases using motor evoked potentials]

[Article in Spanish]

Gomez-Fernandez L, Macias-Gonzalez RJ, Mustelier R, Estrada R, Gamez-Morales L, Paz-Sendin L.

Laboratorio de Neurofisiologia Clinica, Centro Internacional de Restauracion Neurologica, Ciudad de la Habana, Cuba.

INTRODUCTION: The introduction of motor evoked potentials (MEP) into neurological practice has been very useful in the study of motor disorders, especially in patients with motor neurone diseases (MND) and multiple sclerosis (MS). **MATERIAL AND METHODS:** Forty patients with MS, twenty with MND and sixty-four apparently normal control were studied, to evaluate the use of MEP in such patients. Bilateral recordings were made from the abductor pollicis brevis and tibialis anterior. The variables studied were: cortical latency, radicular latency, central conduction time, motor excitation threshold coefficients of amplitude, duration and number of phases. **RESULTS:** There were significant differences in most variables when the control group and the patients were compared, and also between the two subgroups of patients (Wilk's lambda = 0.4197; $p < 0.05$). CA analysis significantly increased the sensitivity of the study (90.4%) with respect to analysis of latencies and central conduction times (70.4%). When analysis of the coefficients of duration and of phases was included, specificity increased by approximately 30% in both groups of patients. **CONCLUSIONS:** The great sensitivity of MEP as a neurophysiological diagnostic tool in investigation of corticospinal bundle lesions was demonstrated.

PMID: 9064163 [PubMed - indexed for MEDLINE]

Abstract Show: 20 Sort

FEB. 11. 2005 11:43AM

AVENTIS US PAT DEPT

NO. 2170 P. 49² OF 2

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Feb 7 2005 09:57:23